

Important Medication Advice

Aspirin, aspirin – related products, ALL herbal products, arthritis and pain medications and Vitamin E must not be taken during the two (2) weeks prior to and after surgery because they increase bleeding. For this reason, it is very important that contents of any "over-the-counter" preparations are checked carefully prior to their use. Many headache preparations, cold remedies and "hangover cures" contain ASPIRIN. The chemical name of aspirin is acetylsalicylic acid. Many arthritis and pain medications have effects on bleeding like aspirin. However, the effects are shorter lived.

TAKING ANY OF THESE MEDICATIONS THE 2 WEEKS PRIOR TO YOUR SURGERY COULD POSSIBLY POSTPONE AND/OR CANCEL YOUR SURGERY BECAUSE OF THE RISK OF BLEEDING

Prescription Medications – Always consults with your physician that prescribed the medication before you stop taking it.

<u>ANTICOAGULANTS</u>

| Acenocoumarin (nicoumalone) | Dicoumarol | Lovenox (enoxaparin) |
|-----------------------------|----------------------|-----------------------|
| Acova (argatroban) | Fragmin (dalteparin) | Miradon (anisindione) |
| Agniomax (bivalirudin) | Heparin | Ticlopidine |
| Coumadin (warfarin) | Pradaxa(dabigatran) | Eliquis (apoxaban) |

Xarelto (rivaroxaban)

ANTIPLATELET DRUGS

| Aggrastat (tirofiban) | Flolan (epoprostenol) | Pletal (cilostazol) |
|-----------------------|---------------------------|----------------------|
| Aggrenox | Integrilin (eptifibtide) | Repro (ebciximab) |
| Agylin (anagrelide) | Plavix (clopidogrel) | St. Joseph 81 mg |
| Aspirin | Persantine (dipyridamole) | Ticlid (ticlopidine) |

PRESCRIPTION PRODUCTS CONTAINING ASPIRIN, SALICYLATES AND/OR SALICYAMIDES

| Forte tabs | Oxycodone |
|----------------|--|
| Inhistone | Pabalate-SF tabs |
| Lanorinal tabs | Pamparin |
| Magan tab | Pepto-Bismol |
| Magsal tabs | Percodan |
| Mammal tabs | Persistin |
| | Lanorinal tabs Magan tab Magsal tabs |

Duragesic Micrainin tabs Robaxisal tabs Easpirin Norgesic Synalgos OC caps Empirin w/codeine tabs Soma Compound tab Nytol Equagesic tabs Nyquil Talwin Compound tab Fiorinal Tabs Oarvon w/ASA pulvules Trisilate tab/liquid Fliorinal w/codeine Oarvon Compound Pulvules Zorprin tabs

NON-PRESCRIPTION DRUGS CONTAINING ASPIRIN, SALICYLATES AND/OR SALICYLAMIDE

4-Way cold tabletsBuf TabsMidol CapletsAlka-SeltzerBuffaprin tabsMobigesic tabAlka-Seltzer Plus coldBufferin tabs (any kind)Momentum tabBuffets II tabsNeocylate tabAnacin tab

Buffinol tab Pabalate Anodynoa tab Buffinol

Extra tab Pepto Bismol Arthralge tab
Cama Arthritis Strength Persistin tab Arthropan liquid

S.A.G. tab Arthritis Strength Bufferin Cope Tab

St. Joseph Childrens aspirin Cosprin Tab St. Joseph Cold tab

A.S.A. Enseals CP 2 tabs Salacol tab Ascriptin tab Dasin Capsule Salato tab

Ascriptin A/D tab Doan's pills Stanback powder

Ascriptin extra strength Duradyn tab Supac

Ecotrin tab Synalgos cap Asperbuf tab Emagrin tab Theracin Aspergum

Empirin tab

Excedrin tab or capsule

Vanquish capsule

Trigesic

Uracel

Bayer Aspirin tab

Goody's powder

Meastruin tab

Verin tab

PRESCRIPTION AND NON-PRESCRIPTION PRODUCTS CONTAINING IBUPROFEN

Always consults with your physician that prescribed the medication before you stop taking it.

Aches-N-Pain Fenoprofen Medipren

Proxicam Advil Fenoprofen calcium

Relafen Aleve Mefanamic Acid Flurbiprofen Mobic Rufen Midol 200 Anaprox Genpril Salflex Anaprox OS Haltran Motrin Solindac Ansaid Ibuprin Nalfon Theracin Antrhotec Indameth Naprosyn Tolectin Clinoril Indocin Naproxen Tolmetin sodium Celebrex Ketoprofen Nuprin Toradol

Daypro Ketorolac Oiclofenac sodium

Trendar Dolobid Lodine Orudis Voltaren Disalid Tromethamine Pamprin IB Dristan Meclofenamate Pediaprofen Feldene

Ponstel Meclomen

HERBAL MEDICATIONS AND NUTRITIONAL SUPPLEMENTS

Horse chestnut bark Sweet birch bark Billberry

Black Cohosh rhizome Kava Kava Sweet Vernal grass leaves Kava kava Licorice root Sweet-scented bedstraw

Niacin Tonka bean seeds Cayenne

Dong quai Ma huang (ephedra) Valerian

Echinacea Meadowsweet flower Vanilla leaf leaves

Feverfew Melatonin Vitamen E Fish oil caps Omega III Willow bark

Garlic Poplar bark or buds Wintergreen leaves Ginseng Red Clover Woodruff plant Ginkgo biloba Yohimbe

Sweet clover plant

Hawthorne St. John's Wort

The following medications MUST BE DISCONTINUED 2 WEEKS before surgery – ONLY with your prescribing doctor's approval. It is very important we know if you are taking these medications

Adipex Corgard Fenphen Inderide Blocadren Diamon Hismanal Levatol Catrol Esidri Inderal Propanalol

Pondimin Redux

NICOTINE interferes with healing by reducing blood flow. You are advised to avoid smoking, nicotine containing gums, and patches 2 weeks prior to and after your surgery. ABSOLUTELY NO SMOKING THE MORNING OF SURGERY!

GLP-1 RECEPTOR AGONISTS

Notify the doctor i are currently taking a GLP-1 receptor agonist (such as Ozempic, Wegovy, Trulicity, Mounjaro, Saxenda, Rybelsus, or Victoza). These medications affect how your stomach empties and may increase the risk of complications during anesthesia or sedation.

To ensure your safety during your upcoming procedure, please follow these instructions carefully:

If you take your GLP-1 medication DAILY:

• Stop taking your medication the day of your procedure.

- Do not take your GLP-1 medication on the morning of the procedure.
- Resume your medication after the procedure only when you are eating normally and per your provider's guidance.

If you take your GLP-1 medication WEEKLY:

- Stop taking your medication 7 days (1 week) before your scheduled procedure.
- Do not resume your medication until after the procedure and only when you are eating normally and your provider advises it is safe to do so.

THESE ARE SOME RECOMMENDATIONS TO HELP PROMOTE HEALING:

- Vitamin C 500 mg twice daily for 1-2 weeks before and after surgery.
- Arnica Montana, take one following surgery and continue for 1-2 weeks after surgery
- Bromelain, begin taking 3 days prior to surgery and continue to 1-2 weeks after surgery

The Arnica Montana and Bromelain can be purchased at your local drug store.

If you have any questions about your medications, you may contact your prescribing physician or call our office.